



Laboratoire d'Économie Forestière



arbre  
Research Institute on the Biology of Trees and Forest Ecosystems



INSTITUT DES SCIENCES ET INDUSTRIES DU VIVANT ET DE L'ENVIRONNEMENT  
PARIS INSTITUTE OF TECHNOLOGY FOR LIFE, FOOD AND ENVIRONMENTAL SCIENCES

# (Forest) Land use impact on water quality

Serge Garcia (LEF-INRA, Nancy)

Impact of the location of agricultural production on  
ecosystem services

Dijon, Monday, November 9th 2015

# Motivations

- **The importance of forest lands in providing ES**
  - The role of the **forest in the supply of a number of goods and services** is recognized by the French policymakers since the Forest Orientation Law (2001)
  - This law reinforces the role of forest policies (with other policies in rural development), **to reduce the greenhouse effect, to preserve biological diversity, to protect soils...**
  - **Water purification** is one of the most important **ecological service provided by (forest) ecosystems** to humanity

# Motivations

- **Effects on land uses on water quality**
  - **Under forest cover, nitrate levels are low**

Land cover	[NO <sub>3</sub> <sup>-</sup> ] in water in the soils at a depth of 1,10m in mg/l
Forests	2
Cut fields	19
Pastures	31
Temporary grassland	28
Winter wheat	46
Rape seed ( <i>colza</i> )	62
Spring cereals	120
Maize as fodder crop ( <i>maïs fourrage</i> )	126

□ Source : Benoit et al. 1997

- **Similar results observed for various pollutants (e.g., pesticides)**

# Motivations

- **The forest has a major positive impact on water quality**
  - **Some wooded formations clearly have a purification role** such as riparian, alluvial forests: the root system has a filtering role and trap nutritive elements (nitrogen, potassium, phosphorus) and some toxic elements
  - **Cultivated lands liberate five times more sediments** into the water course than wooded lands: the forest contributes to protecting land, favoring infiltration, and reducing rapid flow at the surface
  - **Forests generally limit sediment flow and thus turbidity**


# Motivations

- **Influence of forest management on water quality**
  - **Land management in forests is less intensive** than for agriculture and interventions are less frequent
  - **Very limited use of chemicals is made in forest areas:** agro-pharmaceutical products and fertilizers are rarely used
  - **Disturbances in the forest cover**, especially clear cutting, can lead to increase nitrate concentrations in drained water

# Motivations

- **Land uses have an impact on water quality and on costs of drinking water production**
  - **Forest land use is associated with the protection of water resources** from contamination and the **reduced cost of drinking water supply** (Abildtrup and Strange 2000, Willis 2002, Ernst 2004, Freeman et al. 2008)
  - Raw water from catchment areas (for drinking water purpose) with **a large portion of forests is of higher quality than that of agricultural, urban or industrial landscapes**
  - Hence **reducing the need for treatment of drinking water** and, as a result, the associated **costs and prices of drinking water supply**

# Motivations



Drinking water price (2004 ; €/m³ ; IFEN  
SCEES)

Proportion of land area covered by forest  
(2004 ; % ; SCEES)

***Simple correlation or causality?***

# Literature

- Many scientific (mainly hydrological) works on the relationship between forest (and other land uses) and water quality
- A short literature in economics
  - Impacts of alternative land uses on watershed health (Hascic and Wu, 2006; Langpap et al., 2008).
  - Still very few on ***the value of forests in supplying drinking water*** (Núñez et al. 2006, Biao et al. 2010, Elias et al. 2013)
- Vincent et al. (2015) claimed the first econometric analysis on the effect of forests on water treatment costs
  - BUT...



# Literature

- A new approach developed by a LEF team from 2009
  - Objective: estimate the economic value of the ecological service provided by land uses (especially, forest areas) on quality of raw water used for drinking water supply
  - Hypothesis: If land use affects raw water, then we can identify this impact in the process of drinking water supply:  
Raw water is transformed into drinking water with costs
  - Method: estimate a (complete) cost model for drinking water supply by taking the impact of land uses on raw water into account

# Literature


- A new approach from LEF
  - Fiquepron J., Garcia S., Stenger A. (2013). Land use impact on water quality: Valuing forest services in terms of the water supply sector. ***Journal of Environmental Management***, 126, 113-121
  - Abildtrup, J., Garcia S., Stenger A. (2013). The effect of forest land use on the cost of drinking water supply: A spatial econometric analysis. ***Ecological Economics***, 92, 126-136
  - Abildtrup J., Garcia S., Kéré E. (2015). Land use and drinking water supply: A sample selection model with spatial dependence. ***Revue d'Économie Régionale et Urbaine***, 1/2015 (mai), 321-342
  - Abildtrup J., Garcia S., Le Gallo J., Ndiaye Y. (Ongoing work).

# Drinking water supply (DWS)

- DWS covers all operations from resource extraction to customer tap. Several functions with specific costs:
  - Production and treatment: resource extraction (groundwater or surface water) and purification (disinfection, filtering, softening)
  - Transfer and stocking in water tanks and towers and
  - Pressurization of water (gravity or pump-operated system)
  - Distribution to users by distribution mains and service lines
- **Trade-off between treatment costs and distribution costs**, according to water availability and quality
- The sole **analysis of treatment costs could be fake**

# A Production approach...

- A bioeconomic model



## ...Combined with a preference approach

- We estimate welfare changes for the water drinkers
  - We measure the impact of land use changes on water price
  - And determine the value of one ha of afforested land from the water invoice

# Appli 1: Fiquepron et al. (2013)

- System of simultaneous equations

- Quality equations

$$\text{Pesti} = \gamma_0 + \gamma_x X + \gamma_z Z + \varepsilon_{\text{Pesti}}$$

$$\text{NO}_3 = \delta_0 + \delta_x X + \delta_z Z + \varepsilon_{\text{NO}_3}$$

*X : Characteristics of service*

*Z : Land uses*

- Price equation

$$\text{Price} = \alpha_0 + \alpha_x X + \alpha_z Z + \alpha_2 \text{Pesti} + \alpha_3 \text{NO}_3 + \alpha_4 \text{DSP} + \varepsilon_p$$

- Management type equation (DSP)

$$\text{DSP} = \beta_0 + \beta_x X + \beta_z Z + \beta_1 \text{Pesti} + \beta_2 \text{NO}_3 + \varepsilon_d$$

# Empirical strategy

- Econometric analysis

- National study – at the French administrative department level

Sample : 93 administrative departments (without Paris et Corse)

- The econometric estimation method

Estimation of the system of simultaneous equations by GMM

# Results

- Estimation results

Variable to be explained	Explanatory Variable	Impact
<b>Pesticides</b>	% Forest land use	- - -
	% permanent grassland	- - -
	% cereal, oilseeds, protein crops	+ + +
	% vine, arboriculture, market gardening	+ + +
	% underground resources	- - -
<b>Nitrates</b>	% Forest land use	- - -
	% permanent grassland	- - -
	% cereal, oilseeds, protein crops	+ + +
	% vine, arboriculture, market gardening	- - -
	% mountain area	- - -
	Number of pigs per ha	+ + +
	% underground resources	+ + +



# Results

- Estimation results

Variable to be explained	Explanatory Variable	Impact
<b>Management type</b> (% private management)	Delivered drinking water volume	+ + +
	Population density	+ +
	Water recharge	- - -
	Maximal Population	+ + +
	Pesticides	+ +
<b>Drinking water price</b>	Length of water network	+ + +
	% groundwater	- - -
	Management type	+ + +
	Nitrates	+ +

# Results

- Simulation of land use changes

Scenario : substitution forests → mains crops

Change of land uses	Variation / total surface	Surface
% forests	+ 5pts (from 28% to 33%)	+ 2675901 ha
% crops	- 5pts (from 32,5% to 27,5%)	- 2675901 ha

<b>Nitrates</b>	- 2,5 mg/l
<b>Pesticides</b>	- 3,7 pts of outflows to treat

<b>Price</b>	- 0,02 €/m <sup>3</sup>
--------------	-------------------------

<b>Savings on water bill for domestic users</b>	77 million € per year
	<b>29 €/ha/year</b>

## Appli 2: Abildtrup et al. (2013)

- (Positive or negative) externalities between WSS arising from characteristics of neighborhood WSS
  - The costs of water supply may be influenced by **local competition for scarce local water resources** :

If quantity and quality are low, the WSS can decide to use other water resources further (even in another WSS), consequently increasing scarcity and water supply costs for neighboring WSS

- **Technical spillovers** may occur as a result of knowledge diffusion or the sharing of input factors:

and have an impact on the organization of WSS: grouping of municipalities, extension of delegation

# Spatial issues

- **Spatial heterogeneity** can be explained by the proximity of WSS that exhibit the same conditions of operation due to geographical, topographical factors or even the extraction of water in the same aquifer
- **WSS area and land uses generally do not match.** The impact of land uses on service costs should thus be measured by taking both land distribution on the WSS and on its neighbors into account.

# Economic model

- **The costs of supplying drinking water to users can be described by the cost function:**

$$C_i = C(W_i, q_i, X_i, \varepsilon_{ci}; W_{n(j)}, q_{n(j)}, X_{n(j)}, \varepsilon_{cn(j)})$$

- The service  $i$ , the neighboring services  $n(j)$
- $Y$ : delivered water volume,  $X$ : characteristics of the WSS
- $q$ : quality of raw water supposed to dependant on land uses  $L$ :

$$q(L_i, \varepsilon_{qi}; L_{n(j)}, \varepsilon_{qn(j)})$$


- **The pricing model is:**

$$P = AC(W_i, L_i, X_i, \varepsilon_i; W_{n(j)}, L_{n(j)}, X_{n(j)}, \varepsilon_{n(j)}) + R$$

- $AC$ : average cost of drinking water supply
- $P$ : water price,  $R$ : private operator rent

# Data

Sample: 515 communes, which form 283 WSS with 1070 water intakes




- 51 WSS depending on another water basin have missing information
- 232 WSS in the sample
- 21 WSS privately operated (only 9%)

# Data

- PRICE: average drinking water price (composed of a fixed charge and a marginal price) for a consumption of 120 m<sup>3</sup>, without VAT
- MUNICIPAL: number of municipalities served by the WSS
- USER: number of users served by the WSS
- Number of intakes (INTAKE) and type: DRILL (deep), WELL (less), SOURCE (surface)
- ALT-DIF: Difference of altitudes between WSS and intake
- Land uses (in %): FOREST, AGRI, URBAN, OTHER  
Total = 100%
- DELEG = 1 if privately operated




# Spatial distribution of water prices





# Spatial distribution of forest



# Spatial econometric model

- $W$  : a spatial weights matrix to define the dependence between the individual observations
  - Based on neighbourhood contiguity, distance between centroids

- Several spatial econometric models:

$$P = X\beta + L\gamma + W_1Z\delta + \varepsilon$$

with:

$$\varepsilon = \lambda W_2\varepsilon + u$$

- Econometric method (Fingleton and Le Gallo, 2008):
  - 2SLS (multi-step) estimation method

# Estimation results

Variable	Estimate	Estimate
	(Std. Error)	(Std. Error)
	2SLS	Spatial 2SLS
(Intercept)	1.8843*	2.8562**
WATER (in log)	-0.5471**	-0.4708**
USER (in log)	0.6091**	0.4441**
DENSITY (in log)	-0.1978***	-0.1118*
DELEG	0.4866***	0.4510***
MUNICIP	0.0220	0.0216*
DELEG x MUNICIPAL	-0.0241	-0.0268*
INTAKE	0.0070	0.0084
FOREST	-0.9502***	-0.7059***
URBAN	0.0763	-0.0268
OTHER	-0.8201**	-0.6529**
FOREST_lag	--	-0.8405*
URBAN_lag	--	8.1980*
OTHER_lag	--	-1.3144
WATER_lag (in log)	--	0.0489*
$\lambda$	--	0.2557**

Note: \*\*\*: significant at 1%, \*\*: at 5%, \*: at 10%.

# Estimation results

Variable	Estimate (Std. Error)
Spatial 2SLS	
(Intercept)	1.7141*
WATER (in log)	-0.3707**
USER (in log)	0.3504**
DENSITY (in log)	-0.1020*
DELEG	0.4307***
MUNICIP	0.0216*
DELEG x MUNICIPAL	-0.0231
INTAKE	0.0078
FOREST	0.0202
URBAN	0.5567
AGRI	0.6871**
FOREST_lag	-1.1049**
URBAN_lag	9.6119*
AGRI_lag	-1.2841
WATER_lag (in log)	0.0399*

Note: \*\*\*: significant at 1%, \*\*: at 5%, \*: at 10%.

# Estimation results

Average value estimates of forests in supplying water for human consumption (in €)

	No spatial	AGRI as reference	OTHER as reference
Direct impact	-0.0095	-0.0071 <sup>a</sup>	0 <sup>b</sup>
Indirect impact	--	-0.0084	-0.0110
Total impact	-0.0095	-0.0155	-0.0110
Direct value of 1ha	85.08	63.20	0
Indirect value of 1ha	--	75.25	98.93
Total value of 1ha	85.08	138.46	98.93

Note: Values are computed for an average WSS: average water price = €1.08 per m3, average delivered drinking water volume = 104,676 m3, average WSS area = 2165 ha, average proportion of forest lands.

<sup>a</sup> From estimation results, an increase of one point in the proportion of forest (for an equivalent decrease of the proportion of agricultural lands) would imply a decrease of €0.0071 of the water price per m3.

<sup>b</sup> Non significant value.

# Conclusions

- A valuation method for water quality service of forests
- A significant impact of forest land use on water costs
- And a value of the ecological service provided by forests significantly different from zero
- A (relative) value of service different according to different land uses
- Importance to deal with the complete costs of DWS
- Importance of taking spatial issues into account:
  - Forest lands and water supply areas do not match together
  - Spatial spillovers from the water market valuation method

# References

- Abildtrup J., N. Strange (1999). The option value of non-contaminated forest watersheds. *Forest Policy and Economics*, 1, 115–125
- Biao Z., L. Wenhua, W. Gaodi, X. Yu (2010). Water conservation of forest ecosystem in Beijing and its value. *Ecological Economics*, 69(7), 1416–1426
- Elias E., D. Laband, M. Dougherty (2013). Estimating the public water supply protection value of forests. *Journal of Contemporary Water Research & Education*, 152, 94-104
- Ernst C., R. Gullick, K. Nixon (2004). Protecting the source – conserving forests to protect water. *American Water Works Association*, 30(5), 3–7
- Fingleton B., J. Le Gallo (2008). Estimating spatial models with endogenous variables, a spatial lag and spatially dependent disturbances: Finite sample properties. *Papers in Regional Science*, 87(3), 319-339

# References

- Freeman J., R. Madsen, K. Hart, P. Barten, P. Gregory, D. Reckhow, W. Duncan (2008). Statistical analysis of drinking water treatment plant costs, source water quality, and land cover characteristics. Trust for Public Land, San Francisco
- Hascic I., J.J. Wu (2006). Land use and watershed health in the United States. *Land Economics*, 82 (2), 214–239
- Langpap C., I. Hascic, J.J. Wu (2008). Protecting watershed ecosystems through targeted local land use policies. *American Journal of Agricultural Economics*, 90(3), 684–700
- Nuñez D., L. Nahuelhual, C. Oyarzun (2006). Forest and water: the value of native temperate forests in supplying water for human consumption. *Ecological Economics*, 58(3), 606–616



# References

- Vincent J.R., • I. Ahmad, • N. Adnan, W.B. Burwell, S.K. Pattanayak, • J.-S. Tan-Soo, • K. Thomas (2005). Valuing Water Purification by Forests: An Analysis of Malaysian Panel Data. Environmental and Resource Economics, in Press.
- Willis K. (2002). Benefits and Costs of Forests to Water Supply and Water Quality. Report to the Forestry Commission Forestry Commission, Edinburgh



Laboratoire d'Économie Forestière

# Merci

**Contact :** [serge.garcia@nancy.inra.fr](mailto:serge.garcia@nancy.inra.fr)

**Site :** <http://www6.nancy.inra.fr/lef>

**Adresse postale :**

Laboratoire d'Economie Forestière

UMR AgroParisTech/INRA

14 rue Girardet, CS 4216

54042 NANCY Cedex, FRANCE

**Telephone:** +33 3 83 39 68 69